

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/00, 31/55, 31/495, 31/44, 31/47, 31/445		A1	(11) International Publication Number: WO 98/18481
			(43) International Publication Date: 7 May 1998 (07.05.98)
(21) International Application Number: PCT/US97/19880			(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 30 October 1997 (30.10.97)			
(30) Priority Data: 60/029,233 31 October 1996 (31.10.96) US 9711042.3 30 May 1997 (30.05.97) GB			
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SMITH, Roy, G. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). CASCIERI, Margaret, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MACINTYRE, Euan [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MACNEIL, Douglas, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MENKE, John, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			
(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			Published With international search report.
(54) Title: COMBINATION THERAPY FOR THE TREATMENT OF DIABETES AND OBESITY			
(57) Abstract <p>The combination of a metabolic rate modifying agent (e.g., a β_3 adrenergic receptor agonist) and a feeding behavior modifying agent (e.g., a NPY5 antagonist) is useful in the treatment of obesity and diabetes, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients. Methods of treating obesity and diabetes are also described.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

TITLE OF THE INVENTIONCOMBINATION THERAPY FOR THE TREATMENT OF DIABETES
AND OBESITY5 FIELD OF THE INVENTION

The present invention provides a combination useful in the treatment of obesity and diabetes, either as compounds, pharmaceutically acceptable salts or pharmaceutical composition ingredients. Methods of treating obesity and diabetes are also
10 disclosed. More particularly, the combination of the present invention comprises a metabolic rate modifying agent and a feeding behavior modifying agent; and the pharmaceutically acceptable salts and esters thereof.

15 BACKGROUND OF THE INVENTION

Obesity, which can be defined as a body weight more than 20% above the ideal body weight, is a major health concern in Western societies, since it is accompanied by numerous complications such as hypertension, non-insulin dependent diabetes mellitus and
20 arteriosclerosis, which in turn cause heart disease, stroke and premature death. Obesity is the result of a positive energy balance, as a consequence of increased ratio of caloric intake to energy expenditure. The molecular factors regulating food intake and body weight balance are incompletely understood. [B. Staels et al., *J. Biol. Chem.* 270(27),
25 15958 (1995); F. Lonnquist et al., *Nature Medicine* 1(9), 950 (1995)]. Although the genetic and/or environmental factors leading to obesity are poorly understood, several genetic factors have recently been identified.

β -Adrenoceptors have been subclassified as β_1 and β_2 since 1967. Increased heart rate is the primary consequence of β_1 -receptor
30 stimulation, while bronchodilation and smooth muscle relaxation typically result from β_2 stimulation. Adipocyte lipolysis was initially thought to be solely a β_1 -mediated process. However, more recent results indicate that the receptor-mediating lipolysis is atypical in nature. These atypical receptors, later called β_3 -adrenoceptors, are

- 2 -

found on the cell surface of both white and brown adipocytes where their stimulation promotes both lipolysis (breakdown of fat) and energy expenditure

5 Early developments in this area produced compounds with greater agonist activity for the stimulation of lipolysis ($\beta 3$ activity) than for stimulation of atrial rate ($\beta 1$) and tracheal relaxation ($\beta 2$). These early developments disclosed in Ainsworth *et al.*, U.S. Patents 4,478,849 and 4,396,627, were derivatives of phenylethanolamines.


10 Such selectivity for $\beta 3$ -adrenoceptors could make compounds of this type potentially useful as antiobesity agents. In addition, these compounds have been reported to show antihyperglycemic effects in animal models of non-insulin-dependent diabetes mellitus.

15 Recently, assays have been developed which more accurately predict the effects that can be expected in humans. These assays utilize cloned human $\beta 3$ receptors which have been expressed in Chinese hamster ovary cells. See Emorine *et al.*, *Science*, 1989, 245:1118-1121; and Liggett, *Mol. Pharmacol.*, 1992, 42:634-637. The agonist and antagonist effects of the various compounds on the
20 cultivated cells provide an indication of the antiobesity and antidiabetic effects of the compounds in humans.

These developments have recently led to the discovery of potent and selective $\beta 3$ agonists useful for treating obesity and diabetes. For example, U.S. Patent No. 5,451,677, issued September 19, 1995,
25 hereby incorporated by reference, describes substituted phenyl sulfonamides which are selective $\beta 3$ agonists useful for treating obesity and diabetes. These phenyl sulfonamide compounds have been found to be useful in the composition and methods of the instant invention.

30 More recently, a potent and selective $\beta 3$ agonist, (R)-N-[4-[2-[[2-Hydroxy-2-(pyridin-3-yl)ethyl]amino]ethyl]-phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]benzenesulfonamide, hereinafter referred to as Compound A, has been identified.

- 3 -



Compound A

The synthesis of Compound A and its utility for treating obesity and diabetes is described in detail in PCT International application publication number WO 95/29159, published November 2, 1995, and in
5 U.S. Patent No. 5,561,142, issued October 1, 1996. Thus, β_3 agonists, such as Compound A, are useful agents for increasing metabolic rate in mammals.

In addition to β_3 agonists which act on obesity and diabetes by increasing metabolic rate, researchers have recently cloned the
10 mouse OB gene and its human homologue. [Y. Zhang et al., *Nature* 372, 425 (1994)] The OB gene product, i.e., the OB protein (also known as leptin), a 167 amino acid polypeptide, has been shown to result in a dose- and time-dependent weight loss when administered to mice via intraperitoneal (IP) injection. [M.A. Pelleymounter et al.,
15 *Science* 269, 540 (1995)]. This weight loss effect is attributable to both a reduction in food intake and an increase in energy expenditure. Moreover, since both the mouse and human OB protein have this same effect when administered to mice, the possibility exists that similar effects would also occur in humans. [J.L. Halaas et al., *Science* 269, 543
20 (1995)].

Neuropeptide Y (NPY), a 36 amino acid member of the pancreatic polypeptide family with widespread distribution throughout the mammalian nervous system, is another agent which has been identified as being connected with feeding behavior. Neuropeptide Y is
25 involved in regulating eating behavior and is an extremely potent

- 4 -

orixigenic agent [See e.g., Stanley, B.G., et al., *Peptides* 13: 581-587 (1992); Sahu, A. and S.P. Kalra, *Trends In Endocrinology And Metabolism* 4(7): 217-224 (1993)]. When administered intracerebroventricularly or injected into the hypothalamic paraventricular nucleus (PVN) it elicits eating in satiated rats [Clark, J.T., et al., *Endocrinology* 115(1): 427-429 (1984); Stanley, B.G. and S.F. Leibowitz, *Proc. Natl. Acad. Sci. USA* 82: 3940-3943 (1985); Stanley, B.G. and S.F. Leibowitz, *Life Sci.* 35(26): 2635-42 (1984)] and intraventricular injection of antisera to NPY decreases eating [Stanley, B.G., et al., *Peptides*, supra; Sahu, A. and S.P. Kalra, supra]. It has been shown to stimulate appetite in a variety of species and at different stages of development [Stanley, B.G., *Neuropeptide Y in multiple hypothalamic sites controls eating behavior, endocrine, and autonomic systems for body energy balance*, in *Neuropeptide Y*, W.F. Colmers and C. Wahlestedt, Editor. 1993, Humana Press: Totowa, NJ. p. 457-509.]

NPY elicits a broad range of physiologic effects through activation of at least five receptor subtypes known as Y1 [WO 93/09227, published May 13, 1993], Y2 [U.S. Patent No. 5,545,549, issued August 13, 1996; WO 95/21245, published August 10, 1995], Y3, Y4 [U.S. Patent No. 5,516,653, issued May 14, 1996; WO 95/17906, published July 6, 1995] and the "atypical Y1," also known as Y5. Although it has recently been reported that the Y5 receptor is the key subtype responsible for the feeding behavior response in mammals [WO 96/16542, published June 6, 1996], it has now been found that the other subtypes (e.g., NPY1, NPY4) may also be involved in weight control, for example by effects on metabolic rate.

The hypothalamus plays a central role in the integrated regulation of energy homeostasis and body weight, and a number of hypothalamic neuropeptides, for example NPY, galanin, corticotrophin releasing factor (CRF), have been implicated in the mediation of these effects. Additionally, when melanin-concentrating hormone (MCH) was injected into the lateral ventricles of rats, their food consumption increased suggesting that MCH participates in the hypothalamic regulation of feeding behavior; this increase in food consumption was

- 5 -

similar to that seen after galanin administration and somewhat less than seen after NPY treatment. [D. Qu et al., *Nature* **380**: 243-247 (1996)]. Similarly, intracerebroventricular (ICV) glucagon-like peptide 1 (GLP-1) powerfully inhibited feeding in fasted rats suggesting that central
5 GLP-1 is yet another hypothalamic physiological mediator of satiety. [M.D. Turton et al., *Nature* **379**: 69-72 (1996)].

Other mechanisms and agents which have been implicated as playing a role in the regulation of feeding behavior include serotonin reuptake inhibitors such as dexfenfluramine, urocortin agonists, CCK
10 agonists, 5-HT_{2A} agonists and 5-HT_{2C} serotonin receptors. It has been shown that mice lacking 5-HT_{2C} serotonin receptors are overweight as a result of abnormal control of feeding behavior, thereby establishing a role for this receptor in the serotonergic control of appetite. [L.H. Tecott et al., *Nature* **374**: 542-546 (1995)]. A 5-HT_{2C} agonists would
15 therefore be useful for inhibiting food intake. Indeed, a metabolite of dexfenfluramine, (+)norfenfluramine is an agonist at 5-HT_{2C} receptors [M. Spedding et al., *Nature* **380**: 488 (1996)]

It is an object of the present invention to identify compositions useful for the treatment of obesity and/or diabetes. It is a
20 further object of the invention to identify methods of treating obesity or diabetes.

It has now been found that a combination of a metabolic rate modifying agent and an agent which modifies feeding behavior, for example, by reducing total food intake or by reducing caloric intake or
25 selectively reducing intake of specific components of the diet such as carbohydrates or fats, provides effective therapy for treating obesity and diabetes. More specifically, a combination of a β 3 agonist and a NPY5 antagonist is particularly preferred for the treatment of obesity and diabetes.

30

SUMMARY OF THE INVENTION

The present invention provides a composition comprising a metabolic rate modifying agent and a feeding behavior modifying agent; and the pharmaceutically acceptable salts and esters thereof; provided

- 6 -

that when the metabolic rate modifying agent is a β_3 agonist, then the feeding behavior modifying agent is not leptin, or a derivative thereof; and provided further that the metabolic rate modifying agent and the feeding behavior modifying agent are not both the same compound (i.e.,
5 the metabolic rate modifying agent and the feeding behavior modifying agent are not both: leptin, or a derivative thereof, Sibutramine or Ergoset; that is, the metabolic rate modifying agent and the feeding behavior modifying agent are two different compounds).

In one embodiment of the invention is the composition
10 wherein the metabolic rate modifying agent is a compound which increases metabolic rate and the feeding behavior modifying agent is a compound which inhibits feeding behavior.

In a class of the invention is the composition wherein:
(a) the metabolic rate modifying agent is selected from a β_3
15 agonist, leptin, or a derivative thereof, a NPY1 antagonist, a NPY4 antagonist, a UCP1, UCP2 or UCP3 activating agent, Ergoset, a CRF agonist, an agent that inhibits the activity of a specific cAMP dependent protein kinase A (PKA) in adipocytes or a selective inhibitor of phosphodiesterase in adipocytes that would result in increased PKA
20 mediated phosphorylation in adipocytes, or Sibutramine; and
(b) the feeding behavior modifying agent is selected from a NPY5 antagonist, leptin or a derivative thereof, a serotonin reuptake inhibitor, a MCH antagonist, a GLP-1 agonist, a 5-HT_{2C} agonist, a 5-HT_{2A} agonist, a galanin antagonist, a CRF agonist, a urocortin agonist, a
25 melanocortin agonist, an enterostatin agonist, a CCK agonist, Cimetidine, a CCK secretagogue, Ergoset or Sibutramine. Preferably, the metabolic rate modifying agent is selected from a β_3 agonist, leptin, or a derivative thereof, a NPY1 antagonist, a NPY4 antagonist or a UCP1, UCP2 or UCP3 activating agent; and the feeding behavior
30 modifying agent is selected from a NPY5 antagonist, leptin or a derivative thereof, a serotonin reuptake inhibitor, a MCH antagonist, a GLP-1 agonist, a 5-HT_{2C} agonist, a 5-HT_{2A} agonist, a galanin antagonist, a CRF agonist, a urocortin agonist, a melanocortin agonist or an enterostatin agonist.

- 7 -

In a subclass of the invention is the composition wherein the feeding behavior modifying agent is a NPY5 antagonist.

Illustrative of the invention is the composition wherein the metabolic rate modifying agent is selected from a β 3 agonist and a

- 5 NPY1 antagonist. Preferably, the metabolic rate modifying agent is a β 3 agonist. Most preferably, the metabolic rate modifying agent is (R)-N-[4-[2-[[2-hydroxy-2-(pyridin-3-yl)ethyl]amino]ethyl]phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]benzenesulfonamide; or a pharmaceutically acceptable salt and ester thereof.

- 10 An illustration of the invention is the composition wherein the metabolic rate modifying agent is a NPY1 antagonist.

Exemplifying the invention is the composition which comprises leptin, or a derivative thereof, and a NPY5 antagonist; and the pharmaceutically acceptable salts and esters thereof.

- 15 An example of the invention is a pharmaceutical composition comprising any of the compositions described above and a pharmaceutically acceptable carrier. Further illustrating the invention is the pharmaceutical composition made by combining a metabolic rate modifying agent, a feeding behavior modifying agent and a
- 20 pharmaceutically acceptable carrier. Further exemplifying the invention is a process for making a pharmaceutical composition which comprises combining a metabolic rate modifying agent, a feeding behavior modifying agent and a pharmaceutically acceptable carrier; provided that when the metabolic rate modifying agent is a β 3 agonist,
- 25 then the feeding behavior modifying agent is not leptin, or a derivative thereof; and provided further that the metabolic rate modifying agent and the feeding behavior modifying agent are not both the same compound.

- More particularly illustrating the invention is a method of
- 30 treating a condition selected from obesity or diabetes in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of any of the compositions described above.

- 8 -

More specifically exemplifying the invention is a method of treating a condition selected from obesity or diabetes in a subject in need thereof which comprises administering to the subject a metabolic rate increasing agent and a feeding behavior inhibiting agent in an amount effective to treat the condition; provided that when the metabolic rate increasing agent is a $\beta 3$ agonist, then the feeding behavior inhibiting agent is not leptin, or a derivative thereof; and provided further that the metabolic rate increasing agent and the feeding behavior inhibiting agent are not both the same compound.

Another illustration of the invention is the method wherein

(a) the metabolic rate increasing agent is selected from a $\beta 3$ agonist, leptin, or a derivative thereof, a NPY1 antagonist, a NPY4 antagonist or a UCP1, UCP2 or UCP3 activating agent; and

(b) the feeding behavior inhibiting agent is selected from a NPY5 antagonist, leptin or a derivative thereof, a serotonin reuptake inhibitor, a MCH antagonist, a GLP-1 agonist, a 5-HT_{2C} agonist, a 5-HT_{2A} agonist, a galanin antagonist, a CRF agonist, a urocortin agonist, or a CCK antagonist, a melanocortin agonist or an enterostatin agonist.

Preferably, the feeding behavior inhibiting agent is a NPY5 antagonist and the metabolic rate increasing agent is selected from $\beta 3$ agonist and a NPY antagonist. More preferably, the metabolic rate increasing agent is a $\beta 3$ agonist. Most preferably, the metabolic rate increasing agent is (R)-N-[4-[2-[[2-hydroxy-2-(pyridin-3-yl)ethyl]amino]ethyl]phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]benzenesulfonamide.

Another example of the invention is the use of a metabolic rate increasing agent and a feeding behavior inhibiting agent in the preparation of a medicament for the treatment of a condition selected from obesity or diabetes which comprises an effective amount of the metabolic rate increasing agent and the feeding behavior inhibiting agent, together or separately; provided that when the metabolic rate increasing agent is a $\beta 3$ agonist, then the feeding behavior inhibiting agent is not leptin, or a derivative thereof; and provided further that the

- 9 -

metabolic rate increasing agent and the feeding behavior inhibiting agent are not both the same compound.

More specifically illustrating the invention is a drug which is useful for treating a condition selected from obesity or diabetes, the effective ingredients of the said drug being a metabolic rate increasing
5 agent and a feeding behavior inhibiting agent; provided that when the metabolic rate increasing agent is a $\beta 3$ agonist, then the feeding behavior inhibiting agent is not leptin, or a derivative thereof; and provided further that the metabolic rate increasing agent and the feeding
10 behavior inhibiting agent are not both the same compound.

DETAILED DESCRIPTION OF THE INVENTION

This invention is concerned with the combination of certain compounds, or pharmaceutically acceptable salts thereof, for
15 the treatment of obesity and diabetes. Obesity and diabetes mellitus are often treated by encouraging patients to lose weight by reducing their food intake and by increasing their metabolic rate. However, agents that reduce feed intake (e.g., leptin, leptin agonists, 5-HT_{2A} agonists, 5-HT_{2C} agonists, serotonin reuptake inhibitors, NPY5
20 antagonists, CCK agonists, GLP-1 agonists, galanin antagonists, glucagon agonists, MCH agonists) often do not have sustained effects on weight reduction. Thus, it has now been found that combination treatment with an agent that reduces feed intake with an agent that increases metabolic rate (e.g., $\beta 3$ selective agonist, NPY1 antagonist,
25 leptin, leptin agonist, NPY4 antagonist) is advantageous over treatment with either agent alone in the treatment of obesity and diabetes.

As used herein, the terms "selective $\beta 3$ agonist" and " $\beta 3$ selective agonist" are synonymous and refer to agonists which are at
30 least ten-fold selective for the $\beta 3$ adrenergic receptor subtype over the $\beta 1$ and $\beta 2$ adrenergic receptor subtypes in humans. Examples of selective $\beta 3$ adrenergic agonists which are useful in the compositions and methods of the present invention the compounds described in U.S. Patent No. 5,541,677 and PCT International patent application

- 10 -

Publication No. WO 95/29159, published November 2, 1995,
especially Compound A.

5 The term "selective," as used herein in reference to an
agonist or antagonist of a specified receptor subtype, refers to an
agonist or antagonist which binds to the specified receptor subtype
with at least ten-fold greater affinity than it binds to other subtypes
of the receptor.

10 The term "subject," as used herein refers to an animal,
preferably a mammal, most preferably a human, who has been the
object of treatment, observation or experiment.

15 The term "therapeutically effective amount" as used
herein means that amount of active compound(s) or pharmaceutical
agent(s) that elicits the biological or medicinal response in a tissue,
system, animal or human that is being sought by a researcher,
veterinarian, medical doctor or other clinician, which includes
alleviation of the symptoms of the disease being treated.

20 The term "diabetes," as used herein, includes both
insulin-dependent diabetes mellitus (i.e., IDDM, also known as type I
diabetes) and non-insulin-dependent diabetes mellitus (i.e., NIDDM,
also known as Type II diabetes).

25 As used herein, the term "composition" is intended to
encompass a product comprising the specified ingredients in the
specified amounts, as well as any product which results, directly or
indirectly, from combination of the specified ingredients in the specified
amounts.

30 The term "CCK agonist," as used herein refers to a natural
or human made compound that binds to the CCK receptor and acts as an
agonist. Examples of CCK agonists useful in the compositions and
methods of the present invention include, but are not limited to, the
compounds disclosed in C.J. Aquino et al., *J. Med. Chem.* 1996, 39,
562-569. Moreover, utilizing the in vitro assay described by Aquino et
al., one of ordinary skill in the art could readily identify CCK agonists
useful in the present invention.

- 11 -

The term "CCK secretagogue," as used herein refers to a natural or human made compound that causes increased CCK secretion and therefore liberates more endogenous CCK to interact with the CCK receptor.

5 Compounds which reduce food intake include the Ob protein (i.e., leptin), as well as leptin agonists. As used herein, the terms "Ob protein," "OB protein" and "ob protein" all refer to the same protein and are also synonymous with the protein referred to as "leptin." The preparation of the mouse Ob protein is described in
10 Examples 1-2 which follow; human Ob protein can be similarly prepared by one of ordinary skill in the art.

The combination of the present invention is defined as follows:

15 A metabolic rate modifying agent and a feeding behavior modifying agent. Preferably, the combination comprises a metabolic rate increasing agent and a feeding behavior inhibiting agent. Most preferably, the combination comprises a selective $\beta 3$ agonist and an NPY Y5 antagonist.

20 Metabolic rate modifying agents are compounds which, when administered to a subject, act to change the metabolic rate of the subject. Preferably, the metabolic rate modifying agent is a compound which increases metabolic rate when administered to the subject. Metabolic rate modifying agents especially suitable for use in the present invention are those agents which increase metabolic
25 rate by at least 5%, preferably 10%, most preferably 20% in 24 hour energy expenditure when administered to the subject.

Metabolic rate modifying agents are routinely evaluated in rodents (see, e.g., Largis, E.E., et. al., *Drug Devel. Res.* **1994**, 32: 69-76; Carroll, M., et. al., *Diabetes* **1985**, 34: 1198-1202), and, even when
30 inactive in rodents, are tested in additional species such as dog and monkey before ultimately being tested in humans (see, e.g., Connacher, A.A., et. al., *Int'l J. Obesity* **1992**, 16: 685-694; Connacher, A.A., et. al., *Am. J. Clin. Nutr.* **1992**, 55: 258S-261S; Connacher, A.A., et. al., *Brit. Med. J.* **1988**, 296: 1217-1220). One

- 12 -

of ordinary skill in the art can readily identify compounds which are suitable for use as metabolic rate modifying agents in the compositions and methods of the present invention.

Examples of metabolic rate modifying agents include, but are not limited to, β_3 agonists, NPY1 antagonists, NPY4 antagonists, leptin, leptin agonists, and uncoupling protein ("UCP") activating agents, specifically UCP1, UCP2 and/or UCP3 activating agents. Although leptin is typically classified as an agent which inhibits food intake, leptin has other effects. For example, leptin increases metabolic rate by unknown pathways and lowers glucose and insulin in diabetic mice. Indeed, supplementing leptin with an agent that either reduces feed intake or increases metabolic rate has potential for improving the activity of leptin alone; for example, certain obese individuals have elevated endogenous leptin levels and are apparently resistant to leptin. To support this notion, it has been shown that while leptin appears to function by decreasing NPY levels in the hypothalamus it has increased activity in mice lacking the NPY gene compared to wild type mice. [Erickson et al. *Nature* 381: 415-418 (1996)] Thus, for purposes of the present invention, leptin is useful as both a metabolic rate modifying agent as well as a feeding behavior modifying agent, provided that when the metabolic rate modifying agent is leptin then the feeding behavior modifying agent is not leptin, and vice versa (i.e., when the feeding behavior modifying agent is leptin, then the metabolic rate modifying agent is not leptin); and further provided that when the metabolic rate modifying agent is a β_3 agonists, then the feeding behavior modifying agent is not leptin.

Selective β_3 agonist compounds and their use for the treatment of obesity and diabetes are disclosed in U.S. Patent No. 5,541,677 and PCT International patent application Publication No. WO 95/29159, published November 2, 1995. Moreover, Compound A, is synthesized as shown in WO 95/29159 and as shown in Example 70 of U.S. Patent No. 5,561,142, issued October 1, 1996, hereby incorporated by reference. Compound A is (R)-N-[4-[2-[[2-

- 13 -

Hydroxy-2-(pyridin-3-yl)ethyl]amino]-ethyl]-phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]benzenesulfonamide, or a pharmaceutically acceptable salt thereof. Preferably, the dihydrochloride salt of Compound A is utilized in the combinations of the present invention.

The Neuropeptide Y1 receptor and methods of identifying ligands which bind to the Y1 receptor are described in WO 93/09227, published May 13, 1993. Moreover, NPY1 antagonists and their use in the diagnosis and treatment of feeding disorders are described in WO 96/14307, published May 17, 1996. The NPY1 antagonist compounds described in WO 96/14307 are preferred as metabolic rate modifying agents in the compositions and methods of the present invention. Additional NPY Y1 antagonist compounds useful as metabolic rate modifying agents are described in US 5,554,621, EP 0 747 356 A1, EP 0747 357 A1, US 5,552,411, WO 96/14307 and WO 96/40660. The Neuropeptide Y4 receptor and methods of identifying ligands which bind to the Y4 receptor are described in U.S. Patent No. 5,516,653, issued May 14, 1996; and in WO 95/17960, published July 6, 1995. Thus, one of ordinary skill in the art could readily identify NPY4 antagonists useful as metabolic rate modifying agents in the compositions and methods of the present invention.

Other metabolic rate modifying agents include Ergoset (Ergo Science), a dopamine agonist currently in clinical studies for both obesity and diabetes, and Sibutramine (Meridia), a monoamine reuptake inhibitor in development by Knoll (BASF), which is at the pre-registration stage (Europe, US) and is the next anti-obesity agent that the FDA will consider for approval. [SPECTRUM Therapy Markets and Emerging Technologies Decision Resources Inc., Anti-Obesity Market, Press Date: July 26, 1996, pages 103-1 to 103-20]. In addition, a selective CRF agonist [M.Egawa et al. *Neuroscience* 34: 771-775 (1990)], an agent that inhibits the activity of a specific cAMP dependent protein kinase A (PKA) in adipocytes or a selective inhibitor of phosphodiesterase in adipocytes that would result in

- 14 -

increased PKA mediated phosphorylation in adipocytes are also useful in the combinations of the present invention as the metabolic rate modifying agent. Additionally, Uncoupling proteins ("UCP"), particularly UCP1, UCP2 and UCP3, have recently been identified as important mediators of thermogenesis [see, Vidal-Puig, A. et al, 5 *Biochem. Biophys. Res. Commun.* 1997, 235(1): 79-82; Boss, O. et al., *FEBS Lett.* 1997, 408(1): 39-42; Gimeno, R.E. et al., *Diabetes* 1997, 46(5): 900-6; Fleury, C. et al, *Nat. Genet.* 1997, 15(3): 269-72; Zhou, Y.T. et al., *Proc. Natl. Acad. Sci. USA* 1997, 94(12): 10 6386-90]. Thus, UCP activating agents, specifically UCP1, UCP2 or UCP3 activating agents are useful as metabolic rate modifying agents in the instant invention. The utility of such metabolic rate modifying agents is supported by experiments with mice in which the RII-beta gene has been deleted that were shown to be resistant to diet induced 15 obesity. [D. E. Cummings et al. *Nature* 382: 622-626 (1996)].

Feeding behavior modifying agents are compounds which, when administered to a subject, act to change the feeding behavior of the subject. Preferably, the feeding behavior modifying agent is a compound which inhibits feeding behavior and/or reduces 20 food intake when administered to the subject. One of ordinary skill in the art can readily identify feeding behavior modifying agents useful in the compositions and methods of the present invention. For example, compounds which inhibit feeding behavior can be evaluated in rodents according to the procedures described in: Pelleymounter, 25 M.A., et. al., *Science* 1995, 269: 540-543; Halaas, J.L., et. al., *Science* 1995, 269: 543-546; DeVos, P., et. al., *J. Biol. Chem.* 1995, 270: 15958-15961. Examples of feeding behavior modifying agents useful in the present invention include, but are not limited to, leptin, or derivatives thereof, leptin agonists, 5-HT_{2A} 30 agonists, 5-HT_{2C} agonists, serotonin reuptake inhibitors (e.g., dexfenfluramine marketed under the brand name Redux by Servier/Interneuron), NPY5 antagonists, CCK agonists (e.g., FPL-15849: Fisons, Rhone-Poulenc Rorer), GLP-1 agonists, MCH agonists, galanin antagonists, glucagon agonists, Cimetidine

- 15 -

(Tagamet, SmithKline Beecham), Ergoset, Subutramine, urocortin agonists [see, M. Spina, et al., *Science*, 273, 13 September 1996, 1561-1564], melanocortin agonists, specifically melanocortin 4 receptor (MC4R) agonists or mixed MC4R/MC3R agonists [see, 5 Huszar, D. et al., *Cell* 1997, 88: 131-141; Fan, W. et al., *Nature* 1996, 385: 165-168] and enterostatin agonists [see, York, D.A. & Lin, L., *Enterostatin: A Peptide Regulator of Fat Ingestion*, in Molecular & Genetic Aspects of Obesity, 281-297 (G.A. Bray and D.H. Ryan eds., 1996) LSU Press, Baton Rouge; Lin, L. et al., 10 *Peptides* 1997, 18(5): 657-661, describing the effect of enterostatin in chronically reducing fat intake and body weight in rats]. Preferably the feeding behavior modifying agent is selected from a NPY5 antagonist or leptin, or a derivative thereof.

The NPY Y5 receptor and methods of identifying 15 ligands which bind to the Y5 receptor are described in detail in WO 96/16542, published June 6, 1996. One of ordinary skill in the art, following the teaching of WO 96/16542, could readily identify NPY5 antagonist compounds useful in the compositions and methods of the present invention. NPY Y5 antagonists and their use in the 20 diagnosis and treatment of feeding disorders are described in WO 97/19682, WO 97/20820, WO 97/20821, WO 97/20822 and WO 97/20823; these NPY Y5 antagonists are particularly preferred as metabolic rate modifying agents in the compositions and methods of the present invention.

25 The Ob protein is obtained by expression of the recently discovered Ob gene. Expression of mouse Ob gene product in *Drosophila* Schneider 2 (S2) cells is described in Example 13, below. Based on the published sequence of the human Ob cDNA [Y. Zhang et al., *Nature* 372, 425-432 (1994); R.V. Considine et al., *J. Clin.* 30 *Invest.* 95, 2986-2988 (1995)], one of ordinary skill in the art could isolate human cDNA and obtain human Ob protein by expression of the human Ob cDNA in *Drosophila* S2 cells according to the protocol of Example 1. Similarly, the Ob protein may also be

- 16 -

expressed in a bacterial expression system such as *E. coli* or a yeast system and purified by one of ordinary skill in the art.

In addition to the Ob protein, compounds which cause increased expression of the Ob protein (for example, glucocorticoids, *see* P. De Vos, *J. Biol. Chem.* **270**(27), 15958-15961 (1995)) are also useful in combination and methods of the present invention for treating obesity and diabetes. Further included within the invention are derivatives of Ob protein in which the protein is truncated to produce a small peptide and/or one or more amino acids are deleted, added, substituted or modified but which derivatives maintain the biological effect on feeding behavior and food intake and/or metabolic rate. Examples of leptin derivatives (e.g., truncated forms of leptin) which are useful in the present invention include U.S. Patent Nos. 5,552,524; 5,552,523; 5,552,522; 5,521,283; and PCT International application publication nos. WO 96/23513; WO 96/23514; WO 96/23515; WO 96/23516; WO 96/23517; WO 96/23518; WO 96/23519; WO 96/23520, all published on August 8, 1996.

The pharmaceutically acceptable salts of the present invention (in the form of water- or oil-soluble or dispersible products) include the conventional non-toxic salts or the quaternary ammonium salts which are formed, e.g., from inorganic or organic acids or bases. Examples of such acid addition salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-

- 17 -

glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups may be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others. Other pharmaceutically acceptable salts include the sulfate salt ethanolate and sulfate salts.

10 The pharmaceutically acceptable salts of the composition of the instant invention include the composition wherein one of the individual components of the combination is in the form of a pharmaceutically acceptable salt, or the composition wherein all of the individual components are in the form of pharmaceutically acceptable salts (wherein the salts for each of the components can be the same or different), or a pharmaceutically acceptable salt of the combined components (i.e., a salt of the composition). In one embodiment of the present invention, the hydrochloride salt of the composition is utilized.

20 The pharmaceutically acceptable esters in the present invention refer to non-toxic esters, preferably the alkyl esters such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl or pentyl esters, of which the methyl ester is preferred. However, other esters such as phenyl-C₁₋₅ alkyl may be employed if desired.

25 Esterification of alcohols, such as Compound A of the present invention, is performed by a variety of conventional procedures, including reacting the alcohol group with the appropriate anhydride, carboxylic acid or acid chloride. These reactions, as well as other methods of esterification of alcohols, are readily apparent to the skilled artisan.

30 Reaction of the alcohol with the appropriate anhydride is carried out in the presence of an acylation catalyst, such as 4-DMAP (4-dimethylaminopyridine, also known as N,N-dimethylaminopyridine), pyridine, or 1,8-bis[dimethylamino]naphthalene.

- 18 -

Reaction of the alcohol with the appropriate carboxylic acid is carried out in the presence of a dehydrating agent and, optionally, an acylation catalyst. The dehydrating agent, which serves to drive the reaction by the removal of water is selected from dicyclohexylcarbodiimide (DCC), 1-[3-dimethylaminopropyl]-3-ethylcarbodiimide (EDC) 5 or other water soluble dehydrating agents.

Alternatively, reaction of the alcohol with appropriate carboxylic acid can also result in esterification, if performed instead in the presence of trifluoroacetic anhydride, and, optionally, pyridine. A 10 further variant is reacting the alcohol with appropriate carboxylic acid in the presence of N,N-carbonyldiimidazole with pyridine.

Reaction of the alcohol with the acid chloride is carried out with an acylation catalyst, such as 4-DMAP or pyridine.

During any of the above methods for forming esters, it 15 may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in 20 Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

In one aspect, the present invention provides a combination of compounds or a pharmaceutically acceptable ester thereof; or a 25 pharmaceutically acceptable salt thereof for use in the treatment of obesity in human or non-human animals.

The present invention further provides a combination of compounds, or a pharmaceutically acceptable ester thereof; or pharmaceutically acceptable salt thereof, for use in the treatment of 30 hyperglycemia (diabetes) in human or non-human animals.

The disease diabetes mellitus is characterized by metabolic defects in production and utilization of glucose which result in the failure to maintain appropriate blood sugar levels. The result of these defects is elevated blood glucose or hyperglycemia. Research on the

- 19 -

treatment of diabetes has centered on attempts to normalize fasting and postprandial blood glucose levels. Treatments have included parenteral administration of exogenous insulin, oral administration of drugs and dietary therapies.

5 Two major forms of diabetes mellitus are now recognized. Type I diabetes, or insulin-dependent diabetes, is the result of an absolute deficiency of insulin, the hormone which regulates glucose utilization. Type II diabetes, or insulin-independent diabetes (i.e., non-insulin-dependent diabetes mellitus), often occurs in the face of normal,
10 or even elevated levels of insulin and appears to be the result of the inability of tissues to respond appropriately to insulin. Most of the Type II diabetics are also obese. The combination of the present invention is useful for treating both Type I and Type II diabetes. The combination is especially effective for treating Type II diabetes.

15 The combination of compounds of the present invention is useful in the treatment of obesity and diabetes. For these purposes, the combinations of the present invention may be administered orally, parenterally (including subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques), by
20 inhalation spray, or rectally, in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles.

 Thus, in accordance with the combination of the present invention there is further provided a method of treating and a
25 pharmaceutical composition for treating obesity and diabetes. The treatment involves administering to a patient in need of such treatment a pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically effective amount of each compound in the combination of the present invention.

30 These pharmaceutical compositions may be in the form of orally-administrable suspensions or tablets; nasal sprays; sterile injectable preparations, for example, as sterile injectable aqueous or oleaginous suspensions or suppositories.

- 20 -

In accordance with the methods of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. For example, in a two-component combination which is the β_3 agonist, Compound A, and a NPY5 antagonist, treatment with the NPY5 antagonist can commence prior to, subsequent to or concurrent with commencement of treatment with Compound A. Furthermore, the term administering also encompasses the use of prodrugs of the β_3 agonist and/or NPY5 antagonist which convert in vivo to a selective β_3 agonist or NPY5 antagonist. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly.

When any of the active ingredients (e.g. Compound A) are administered orally as a suspension, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents known in the art. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

When administered by nasal aerosol or inhalation, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

The compounds utilized in the combination may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or

- 21 -

intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. When administered by injection, the injectable solutions or suspensions may be formulated according to known art, using suitable non-toxic, parenterally-acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

When rectally administered in the form of suppositories, these compositions may be prepared by mixing the drug with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquidify and/or dissolve in the rectal cavity to release the drug.

The active ingredients of the combination (e.g., Compound A) of the present invention may be administered as a pharmaceutical composition, for example, with an inert diluent, or with an assimilable edible carrier, or they may be enclosed in hard or soft shell capsules, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, which includes sublingual administration, these active compounds may be incorporated with excipients and used in the form of tablets, pills, capsules, ampules, sachets, elixirs, suspensions, syrups, and the like. Such compositions and preparations should contain at least 0.1 percent of the active ingredients. The percentage of active ingredients in these compositions may, of course, be varied and may conveniently be between about 2 percent to about 60 percent of the weight of the unit. The amount of active ingredients in such therapeutically useful compositions is such that an effective dosage will be obtained. The active compounds can also be administered intranasally as, for example, liquid drops or spray.

The effective dosage of each of the active ingredients employed in the combination may vary depending on the particular

- 22 -

compound employed, the mode of administration, the condition being treated and the severity of the condition being treated. Thus, the dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound thereof employed. A physician, clinician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentration of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug.

The compounds of this invention can be administered to humans in the dosage ranges specific for each compound. In general, for treating obesity and/or diabetes, the metabolic rate modifying agent is administered at a daily dosage of from about 0.001 to about 20 mg/kg of animal body weight, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form; in the case of an adult human, the total daily dose will generally be from about 0.07 milligrams to about 3500 milligrams of the metabolic rate modifying agent. In general, for treating obesity and/or diabetes, the feeding behavior modifying agent is administered at a daily dosage of from about 0.001 to about 20 mg/kg of animal body weight, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form; in the case of an adult human, the total daily dose will generally be from about 0.07 milligrams to about 3500 milligrams of the feeding behavior modifying agent.

More specifically, when treating diabetes mellitus and/or hyperglycemia generally satisfactory results are obtained when Compound A, or a pharmaceutically acceptable salt thereof, is administered at a daily dosage of from about 0.001 milligram to about 20 milligram per kilogram of animal body weight, preferably given in a

- 23 -

single dose or in divided doses two to six times a day, or in sustained release form. In the case of an adult human, the total daily dose of Compound A will generally be from about 0.07 milligrams to about 3500 milligrams. The Ob protein is administered at a daily dosage of

5 from about 0.05 mg/kg to about 20 mg/kg, preferably injected in a single dose or in divided doses 2 to 3 times per day, or in sustained release form. Preferably, the daily dosage of Ob protein is from about 0.05 mg/kg to about 5 mg/kg. The NPY5 antagonist is administered at a daily dosage of from about 0.001 mg/kg to about

10 20 mg/kg, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form; in the case of an adult human, the total daily dose of NPY5 antagonist will generally be from about 0.07 milligrams to about 3500 milligrams. The NPY1 antagonist is administered at a daily dosage of from about 0.001 mg/kg to about 20

15 mg/kg, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form; in the case of an adult human, the total daily dose of NPY1 antagonist will generally be from about 0.07 milligrams to about 3500 milligrams. The dosage regimen of any of the individual components of the combination may be adjusted to

20 provide the optimal therapeutic response. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body

25 weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

When treating obesity, in conjunction with diabetes and/or hyperglycemia, or alone, generally satisfactory results are obtained

30 when Compound A is administered at a daily dosage of from 0.01 milligram to about 20 milligrams per kilogram of animal body weight, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form. In the case of an adult human, the total daily dose of Compound A will generally be from about 0.7

- 24 -

milligrams to about 3500 milligrams. The Ob protein is administered at a daily dosage of from about 0.05 mg/kg to about 20 mg/kg, preferably given in a single dose or in divided doses 2 to 3 times per day, or as a constant infusion. Preferably, the daily dosage of Ob protein is from

5 about 0.05 mg/kg to about 5 mg/kg. The NPY5 antagonist is administered at a daily dosage of from about 0.001 mg/kg to about 20 mg/kg, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form; in the case of an adult human, the total daily dose of NPY5 antagonist will generally be from about

10 0.07 milligrams to about 3500 milligrams. The NPY1 antagonist is administered at a daily dosage of from about 0.001 mg/kg to about 20 mg/kg, preferably given in a single dose or in divided doses two to six times a day, or in sustained release form; in the case of an adult human, the total daily dose of NPY1 antagonist will generally be from about

15 0.07 milligrams to about 3500 milligrams. The dosage regimen of any of the components of the combination may be adjusted to provide the optimal therapeutic response. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the

20 activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

25 The tablets, pills, capsules, and the like may also contain a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin. When a dosage

30 unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as a fatty oil.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in

- 25 -

addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

These active compounds may also be administered
5 parenterally. Solutions or suspensions of these active compounds can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary
10 conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or
15 dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for
20 example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

The following examples are provided so that the invention might be more fully understood. They should not be construed as
25 limiting the invention in any way.

- 26 -

EXAMPLE 1

Expression of mouse Ob gene product in Drosophila S2 cells

Total white adipose RNA was isolated from Swiss-
5 Webster mice and first strand cDNA synthesized. Using the
polymerase chain reaction (PCR), the coding region of the ob cDNA
was isolated as 2 overlapping fragments using the following primer
sets (5' CAGTGAGCCCCAAGAAGAGG 3' (SEQ. I.D. NO: 1),
5' TCCAGGTCATTGGCTATCTG 3' (SEQ. I.D. NO: 2)) and (5'
10 ATTCCTGGGCTTCAGGGGATTCTGAGTTTC 3' (SEQ. I.D. NO:
3), 5' GCGTGTACCCACGGAGGAAC 3' (SEQ. I.D. NO: 4)). The
resulting 380 bp and 626 bp fragments were purified and used as
templates in a subsequent PCR reaction with primers
5' AAGAATTCATGTTGCTGGAGACCCCTGTGTC 3' (SEQ. I.D.
15 NO: 5) and 5' AAGGATCCTCAGCATTACAGGGCTAACATC 3'
(SEQ. I.D. NO: 6).

The final 501 bp fragment was sequenced using dye-
primer chemistry on a Perkin-Elmer/Applied Biosystems 373A
sequencer. The deduced amino acid sequence encoded by this cDNA
20 was identical to that previously described. [Y. Zhang et al., *Nature*
373, 425-432 (1994)]. The 501 bp fragment encoding the ob gene
product was subcloned via EcoRI and BamHI sites into plasmid
pRmHa3 [T.A. Bunch et al., *Nucleic Acids Research* 16, 1043-1061
(1988)]. Drosophila S2 cells were cotransfected with pUChsneo [H.
25 Steller and V. Pirrotta, *EMBO Journal* 4, 167-171 (1985)] and the
ob expression or control (pRmHa3) plasmid by CaPO₄ precipitation.
A polyclonal population of transfected cells was selected with 1
mg/ml G418 and grown under serum free conditions in EXCELL
401 medium (JRH Bioscience). Cells were seeded at 2 x 10⁶ cells/ml
30 and CuSO₄ added to a final concentration of 1 mM. After 7 days,
cells were removed by centrifugation and the supernatants filtered
through a 0.45 µM filter.

- 27 -

Partial purification of the ob protein:

- Proteins in the supernatant were concentrated by precipitation with 50 percent (NH₄)SO₄. The precipitate was dissolved in Buffer A (20mM Tris pH 8, 1mM DTT and 1mM EDTA) and desalted over PD-10 columns (Pharmacia) equilibrated in Buffer A. Proteins in the PD-10 eluate were subjected to Mono Q chromatography (Pharmacia) and eluted with a 0-200 mM gradient of NaCl in Buffer A. Peak fractions of ob immunoreactivity (centered at 100 mM NaCl) were identified using antiserum 103-2.
- Based on densitometric scanning of polyacrylamide gel containing pooled fractions eluted from the Mono Q column, the ob protein was 30 percent pure after this step. 0.5 to one milligram of this partially purified ob protein preparation was obtained per liter of S2 cells. Peak fractions were analyzed by HPLC electrospray mass spectrometry using a C₄ column (1 x 100 mm) eluted in a linear gradient of acetonitrile (zero to sixty-seven percent in 10mM trifluoroacetic acid. This reverse phase HPLC/mass spectrometry analysis demonstrated the immunoreactive protein to have a molecular mass of 16,004 Daltons, identical to that predicted for the ob gene product after cleavage of the N terminal signal sequence between amino acids 21 and 22.

EXAMPLE 2

Immunological Methods

- Antiserum 103-2 was isolated from a New Zealand white rabbit injected with a 4-branch multiple antigenic peptide corresponding to amino acids 22-41 of the mouse ob sequence, i.e., VPIQKVQDDTKTLIKTIVTR (SEQ. I.D. NO: 7). [Y. Zhang et al., *Nature* 372, 425-432 (1994)]. Western blot analysis was performed on nitrocellulose (BA85m 0.45 µm pore size, Schleicher and Schuell Inc.). Immunodetection of the ob gene product was performed in TBS-T (20 mM Tris-Cl pH 7.6, 137 mM NaCl, 0.1% Tween 20) utilizing the ECL kit (Amersham). The secondary antibody (anti-

- 28 -

rabbit Ig, horseradish peroxidase linked F(ab')₂ fragment, Amersham) was used at 1:3000 dilution. A single immunoreactive protein with an apparent molecular weight of 14.5 kDa was identified.

5

EXAMPLE 3

In vivo study for combination therapy with Compounds B and C

Compound B is defined as a molecule that inhibits feed intake such as a selective serotonin 5HT_{2C} receptor subtype agonist, an NPY₅ selective receptor antagonist, a galanin antagonist, a CCK agonist, an MCH antagonist, a CRF agonist, a urocortin agonist, a glucagon-like peptide agonist, a glucagon agonist, leptin, leptin agonist. Compound C is a compound that increases metabolic rate such as a selective beta-3 adrenergic receptor agonist (e.g., Compound A), an NPY₁ receptor antagonist, leptin, leptin agonist. Human patients are given by injection (i.e., subcutaneous, intramuscular or intravenous) or orally of 0.001 to 100 mg/kg of Compound B, administered one to three times per day, together with or followed by 0.001 to 100 mg/kg Compound C, administered orally or by injection one to three times per day. The first treatment is given on day 0, and the patients are treated daily for a six month period. A set of control patients are untreated (e.g., placebo) for comparison. Body weight data are collected each week and a sustained decrease in body weight of at least 2% is observed. Statistical significance of body weight decreases is determined by performing a two-factor (group and day) analysis of variance (ANOVA) with repeated measures followed by a post hoc Least Squares Difference (LSD) test.

Determination of plasma insulin and glucose levels:

Blood is collected into heparinized capillary tubes on the day before the first treatment and daily for the first week. Thereafter, patients should individually monitor their own blood glucose daily (using commercially available kits). Blood is collected for laboratory analysis weekly at the same time that body weight measurements are

- 29 -

taken. Blood samples are analyzed (e.g., at a registered hospital or other GLP laboratory) for glucose and insulin levels.

- 5 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations, or modifications, as come within the scope of the following claims and its equivalents.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(I) APPLICANT: SMITH, ROY G.
CASCIERI, MARGARET A.
MACINTYRE, EUAN
MACNEIL, DOUGLAS J.
MENKE, JOHN G.

(II) TITLE OF INVENTION: COMBINATION THERAPY FOR THE TREATMENT OF
DIABETES AND OBESITY

(III) NUMBER OF SEQUENCES: 7

(IV) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: MARY A. APPOLLINA - MERCK & CO., INC.
(B) STREET: 126 EAST LINCOLN AVENUE - P.O. BOX 2000
(C) CITY: RAHWAY
(D) STATE: NEW JERSEY
(E) COUNTRY: USA
(F) ZIP: 07065-0907

(V) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: FLOPPY DISK
(B) COMPUTER: IBM PC COMPATIBLE
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PATENTIN RELEASE #1.0, VERSION #1.30

(VI) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US
(B) FILING DATE:
(C) CLASSIFICATION:

(VIII) ATTORNEY/AGENT INFORMATION:

(A) NAME: APPOLLINA, MARY A.
(B) REGISTRATION NUMBER: 34,087
(C) REFERENCE/DOCKET NUMBER: 19822Y

(IX) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (732) 594-3462
(B) TELEFAX: (732) 594-4720

(2) INFORMATION FOR SEQ ID NO:1:

(I) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 BASE PAIRS
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: SINGLE
(D) TOPOLOGY: LINEAR

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAGTGAGCCC CAAGAAGAGG

(2) INFORMATION FOR SEQ ID NO:2:

- (I) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 BASE PAIRS
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 (D) TOPOLOGY: LINEAR

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCCAGGTCAT TGGCTATCTG

20

(2) INFORMATION FOR SEQ ID NO:3:

- (I) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 BASE PAIRS
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 (D) TOPOLOGY: LINEAR

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATTCCTGGGC TTCAGGGGAT TCTGAGTTTC

30

(2) INFORMATION FOR SEQ ID NO:4:

- (I) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 BASE PAIRS
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 (D) TOPOLOGY: LINEAR

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GCGTGTACCC ACGGAGGAAC

20

(2) INFORMATION FOR SEQ ID NO:5:

- (I) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 BASE PAIRS
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 (D) TOPOLOGY: LINEAR

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AAGAATTCAT GTTGCTGGAG ACCCCTGTGT C

31

(2) INFORMATION FOR SEQ ID NO:6:

- (I) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 BASE PAIRS
 (B) TYPE: NUCLEIC ACID

- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:6:

AAGGATCCTC AGCATTTCAGG GCTAACATC

29

(2) INFORMATION FOR SEQ ID NO:7:

- (I) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 AMINO ACIDS
 - (B) TYPE: AMINO ACID
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: LINEAR

(II) MOLECULE TYPE: PROTEIN

(XI) SEQUENCE DESCRIPTION: SEQ ID NO:7:

VAL	PRO	ILE	GLN	LYS	VAL	GLN	ASP	ASP	THR	LYS	THR	LEU	ILE	LYS	THR
1				5					10					15	

ILE	VAL	THR	ARG
			20

- 33 -

WHAT IS CLAIMED IS:

1. A composition comprising a metabolic rate
modifying agent and a feeding behavior modifying agent; and the
5 pharmaceutically acceptable salts and esters thereof; provided that when
the metabolic rate modifying agent is a $\beta 3$ agonist, then the feeding
behavior modifying agent is not leptin, or a derivative thereof; and
provided further that the metabolic rate modifying agent and the
feeding behavior modifying agent are not both the same compound.
10
2. The composition of Claim 1, wherein the metabolic
rate modifying agent is a compound which increases metabolic rate and
the feeding behavior modifying agent is a compound which inhibits
feeding behavior.
15
3. The composition of Claim 2, wherein
(a) the metabolic rate modifying agent is selected from a $\beta 3$
agonist, leptin, or a derivative thereof, a NPY1 antagonist, a NPY4
antagonist or a UCP1, UCP2 or UCP3 activating agent; and
20 (b) the feeding behavior modifying agent is selected from a NPY5
antagonist, leptin, or a derivative thereof, a serotonin reuptake
inhibitor, a MCH antagonist, a GLP-1 agonist, a 5-HT_{2C} agonist, a 5-
HT_{2A} agonist, a galanin antagonist, a CRF agonist, a urocortin agonist,
a melanocortin agonist or an enterostatin agonist.
25
4. The composition of Claim 3, wherein the feeding
behavior modifying agent is a NPY5 antagonist.
5. The composition of Claim 4, wherein the metabolic
30 rate modifying agent is selected from a $\beta 3$ agonist or a NPY1
antagonist.
6. The composition of Claim 5, wherein the metabolic
rate modifying agent is a $\beta 3$ agonist.

- 34 -

7. The composition of Claim 6, wherein the β_3 agonist is (R)-N-[4-[2-[[2-hydroxy-2-(pyridin-3-yl)ethyl]amino]ethyl]phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]benzenesulfonamide; or a pharmaceutically acceptable salt and ester thereof.

5

8. The composition of Claim 5, wherein the metabolic rate modifying agent is a NPY1 antagonist.

9. The composition of Claim 3, which comprises leptin, or a derivative thereof, and a NPY5 antagonist; and the pharmaceutically acceptable salts and esters thereof.

10

10. A pharmaceutical composition comprising the composition of Claim 1 and a pharmaceutically acceptable carrier.

15

11. The pharmaceutical composition of Claim 10, made by combining a metabolic rate modifying agent, a feeding behavior modifying agent and a pharmaceutically acceptable carrier.

20

12. A process for making a pharmaceutical composition which comprises combining a metabolic rate modifying agent, a feeding behavior modifying agent and a pharmaceutically acceptable carrier; provided that when the metabolic rate modifying agent is a β_3 agonist, then the feeding behavior modifying agent is not leptin, or a derivative thereof; and provided further that the metabolic rate modifying agent and the feeding behavior modifying agent are not both the same compound.

25

13. A method of treating obesity in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of the composition of Claim 2.

30

- 35 -

14. A method of treating diabetes in a subject in need thereof which comprises administering to the subject a therapeutically effective amount of the composition of Claim 2.

5 15. A method of treating a condition selected from obesity or diabetes in a subject in need thereof which comprises administering to the subject a metabolic rate increasing agent and a feeding behavior inhibiting agent in an amount effective to treat the condition; provided that when the metabolic rate increasing agent is a
10 β 3 agonist, then the feeding behavior inhibiting agent is not leptin, or a derivative thereof; and provided further that the metabolic rate increasing agent and the feeding behavior inhibiting agent are not both the same compound.

15 16. The method of Claim 15, wherein
 (a) the metabolic rate increasing agent is selected from a β 3 agonist, leptin, or a derivative thereof, a NPY1 antagonist, a NPY4 antagonist or a UCP1, UCP2 or UCP3 activating agent; and
 (b) the feeding behavior inhibiting agent is selected from a NPY5
20 antagonist, leptin or a derivative thereof, a serotonin reuptake inhibitor, a MCH antagonist, a GLP-1 agonist, a 5-HT_{2C} agonist, a 5-HT_{2A} agonist, a galanin antagonist, a CRF agonist, a urocortin agonist, a melanocortin agonist or an enterostatin agonist.

25 17. The method of Claim 16, wherein the feeding behavior inhibiting agent is a NPY5 antagonist.

 18. The method of Claim 17, wherein the metabolic rate increasing agent is a selected from β 3 agonist and a NPY antagonist.

30 19. The method of Claim 18, wherein the metabolic rate increasing agent is a β 3 agonist.

- 36 -

20. The method of Claim 19, wherein the metabolic rate increasing agent is (R)-N-[4-[2-[[2-hydroxy-2-(pyridin-3-yl)ethyl]amino]ethyl]phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]benzenesulfonamide.

5

21. The use of a metabolic rate increasing agent and a feeding behavior inhibiting agent in the preparation of a medicament for the treatment of a condition selected from obesity or diabetes which comprises an effective amount of the metabolic rate increasing agent and the feeding behavior inhibiting agent, together or separately; provided that when the metabolic rate increasing agent is a β_3 agonist, then the feeding behavior inhibiting agent is not leptin, or a derivative thereof; and provided further that the metabolic rate increasing agent and the feeding behavior inhibiting agent are not both the same compound.

10

15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/19880**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 38/00, 31/55, 31/495, 31/44, 31/47, 31/445

US CL :514/12, 218, 252, 278, 290, 307, 318, 337, 338, 340, 342

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/12, 218, 252, 278, 290, 307, 318, 337, 338, 340, 342

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	US 5,635,503 A (POINDEXTER et al.) 03 June 1997, see the abstract, column 5, lines 42-43 and column 6, line 38 - column 7, line 20.	1-5, 8, 8, 10-13, 15 - 18 and 21. NON
Y, P	US 5,602,024 A (GERALD et al.) 11 February 1997, see column 2, lines 5-10 and column 32, lines 55-60.	1-21
Y	US 5,561,142 A (FISHER et al.) 01 October 1996, see the abstract, column 32, line 30 - column 33, line 42 and column 48, Example 70.	1-21
Y	US 5,554,621 A (POINDEXTER et al.) 10 September 1996, see the abstract, column 6, lines 14-15 and column 7, lines 10-59.	1-5, 8, 10-13, 15-18 and 21.



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 JANUARY 1998

Date of mailing of the international search report

06 FEB 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RAYMOND J. HENLEY JR.

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/19880

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,552,524 A (BASINSKI et al.) 03 September 1996, see the abstract, column 2, line 40 - column 4, line 35 and column 16, line 48 - column 17, line 30.	1-21